

# Toronto Western Hospital ANTIBIOGRAM

## All Inpatients and Outpatients

### January 1, 2014 - December 31, 2014

#### Blood Isolates – % Susceptible

<div style="display: flex; align-items: center;"> <div style="margin-right: 10px;"> <p>≥80% Susceptible</p> <p>70-79% Susceptible</p> <p>≤69% Susceptible</p> </div> <div style="display: flex; flex-direction: column; align-items: center;"> <div style="width: 20px; height: 20px; background-color: #90EE90; margin-bottom: 5px;"></div> <div style="width: 20px; height: 20px; background-color: #FFFF00; margin-bottom: 5px;"></div> <div style="width: 20px; height: 20px; background-color: #FF0000;"></div> </div> </div>		#	%	Fluconazole	Voriconazole	Caspofungin	Micafungin	Anidulafungin

<b>ALL CANDIDA</b>	<b>12</b>	<b>108</b>						
<i>Candida albicans</i>	3	25	100	100	100	100	100	100
<i>Candida parapsilosis</i>	3	33	100	100	100	100	100	100
<i>Candida glabrata</i> *	3	25			100	100	100	
<i>Candida guilliermondii</i>	1	8	100	100	100	100	100	100
<i>Candida species</i>	1	8	100	100	100	100	100	100
<i>Candida lusitanae</i> ^	1	8						

#### Year-Specific Notes:

- Interpretive standards for fluconazole, voriconazole, caspofungin, micafungin and anidulafungin susceptibility for all *Candida spp.* were changed in 2013 resulting in an apparent drop in susceptibility to some of these antifungal agents compared to past years.

#### Organism-Specific Notes:

- \* *Candida glabrata* - There are no interpretative breakpoints for voriconazole.  
There are no susceptibility breakpoints for fluconazole, the percentage provided refers to the percentage of *Candida glabrata* that are susceptible dose-dependent.

- ^ *Candida lusitanae* - There are no interpretative breakpoints for any antifungal.

---

Prepared by University Health Network/Mount Sinai Hospital Department of Microbiology

May 28, 2016